The user of this document is responsible to confirm that the Certificate matches the product of interest.



CERTIFICATE OF ANALYSIS

VIAL CONTENTS: Endosafe® Control Standard Endotoxin is prepared from <i>E. coli</i> strain 055:B5. Each vial contains 10 ng of purified Lipopolysaccharide, freeze dried in a stabilized matrix.							
RSE/CSE RATIO: The potency of this standard in Endotoxin units, (EU) has been determined to be 13 EU/ng by the method formerly described in Appendix C (<i>Kinetic Technique</i>) of the GUIDELINE ON VALIDATION OF THE LIMULUS AMEBOCYTE LYSATE TEST AS AN END PRODUCT ENDOTOXIN TEST FOR HUMAN AND ANIMAL PARENTERAL DRUGS, BIOLOGICAL PRODUCTS, AND MEDICAL DEVICES, published by the U.S. Food and Drug Administration.							
CSE Lot:	EX03492	CSE Exp. Date: _	Dec 2023				
LAL Reagent Lot:	M3303L	LAL Exp. Date: _	May 2024				
RSE Lot:	H0K354	RSE/CSE Ratio: _	13	EU/ng	Vial contents: _	130	_ EU/vial
IS/CSE RATIO: The Expert Committee on Biological Standardization of WHO has assigned a potency of the IS as 10,000 IU per vial of IS, so that 1 IU = 1 EU. The potency of this endotoxin standard in International (Endotoxin) Units, IU, has been designated asIU/ng. DIRECTIONS FOR USE: Reconstitute the lyophilized material with26mL of LAL reagent grade water to obtain50EU/mL or50IU/mL. Vortex mix vigorously for at least 5 minutes after rehydration, and for at least 1 minute immediately prior to each use.							
STORAGE: Store rehydrated material at 2-8°C for up to 4 weeks. Store lyophilized material at controlled room temperature or refrigerated as preferred. Diluted endotoxin should not be stored except under validated conditions.							
CAUTION: DO NOT FREEZE ENDOTOXIN SOLUTIONS							
Prepared By:	Jestin	n Jase		Dat	e: 22 M	01/100	и
Quality Review	ed By:(Pourjai		Date	e: 24Mar	<u> </u>	