

CERTIFICATE OF ANALYSIS

VIAL CONTENTS: Endosafe® Control Standard Endotoxin is prepared from *E. coli* strain 055:B5. Each vial contains 500 ng of purified Lipopolysaccharide, freeze dried in a stabilized matrix.

RSE/CSE RATIO: The potency of this standard in Endotoxin units, (EU) has been determined to be	
20 EU/ng by the method formerly described in Appe GUIDELINE ON VALIDATION OF THE LIMULUS AMEBOCYTE LYSAT TEST FOR HUMAN AND ANIMAL PARENTERAL DRUGS, BIOLOGIC published by the U.S Food and Drug Administration.	endix C (Gel-clot Technique) of the E TEST AS AN END PRODUCT ENDOTOXIN
CSE Lot:EM03233	_
LAL Reagent Lot: M3303L LAL Exp. Date: May 2024	_
RSE Lot: <u>H0K354</u> RSE/CSE Ratio: <u>20</u>	_EU/ng Vial contents:10000 EU/vial
Geometric Mean Sensitivity with RSE:EU/mL	
IS/CSE RATIO: The Expert Committee on Biological Standardization of WHO has assigned a potency of the IS as 10,000 IU per vial of IS, so that 1 IU = 1 EU. The potency of this endotoxin standard in International (Endotoxin) Units, IU, has been designated asIU/ng.	
DIRECTIONS FOR USE: Reconstitute the lyophilized material with5.0mL of LAL reagent grade water to obtain2000EU/mL or2000IU/mL. Vortex mix vigorously for at least 5 minutes after rehydration, and for at least 1 minute immediately prior to each use.	
STORAGE: Store rehydrated material at 2-8°C for up to 4 weeks. Store lyophilized material at controlled room temperature or refrigerated as preferred. Diluted endotoxin should not be stored except under validated conditions.	
CAUTION: DO NOT FREEZE ENDOTOXIN SOLUTIONS	
Prepared By: Sauth	Date: 08 Mar 202 (
Quality Reviewed By: Thu E. Thu	Date: 13MAR 2021
	CA-GC500-07