

CERTIFICATE OF ANALYSIS

VIAL CONTENTS: Endosafe® Control Standard Endotoxin is prepared from E. coli strain 055:B5. Each vial contains 500 ng of purified Lipopolysaccharide, freeze dried in a stabilized matrix. RSE/CSE RATIO: The potency of this standard in Endotoxin units, (EU) has been determined to be _EU/ng by the method formerly described in Appendix C (Gel-clot Technique) of the GUIDELINE ON VALIDATION OF THE LIMULUS AMEBOCYTE LYSATE TEST AS AN END PRODUCT ENDOTOXIN TEST FOR HUMAN AND ANIMAL PARENTERAL DRUGS, BIOLOGICAL PRODUCTS, AND MEDICAL DEVICES, published by the U.S Food and Drug Administration. EM84083 CSE Exp. Date: Jan 2022 CSE Lot: L4242L LAL Exp. Date: Jul 2022 LAL Reagent Lot: ___ H0K354 RSE/CSE Ratio: _____ 10 ____ EU/ng Vial contents: ____ 5000 EU/vial RSE Lot: 0.03 EU/mL Geometric Mean Sensitivity with RSE: IS/CSE RATIO: The Expert Committee on Biological Standardization of WHO has assigned a potency of the IS as 10,000 IU per vial of IS, so that 1 IU = 1 EU. The potency of this endotoxin standard in International (Endotoxin) Units, IU, has been designated as _____10 ___IU/ng. DIRECTIONS FOR USE: Reconstitute the lyophilized material with _____5.0 ___mL of LAL reagent grade water to obtain 1000 EU/mL or 1000 IU/mL. Vortex mix vigorously for at least 5 minutes after rehydration, and for at least 1 minute immediately prior to each use. STORAGE: Store rehydrated material at 2-8°C for up to 4 weeks. Store lyophilized material at controlled room temperature or refrigerated as preferred. Diluted endotoxin should not be stored except under validated conditions. **CAUTION: DO NOT FREEZE ENDOTOXIN SOLUTIONS** d By: Jame Date: 22 FEB 2019 ____ Date: 19 Feb 2019 Quality Reviewed By:_

CA-GC500-06