The user of this document is responsible to confirm that the Certificate matches the product of interest.



## **CERTIFICATE OF ANALYSIS**

VIAL CONTENTS: Endosafe® Control Standard Endotoxin is prepared from E. coli strain 055:B5. Each vial

contains 10 ng of purified Lipopolysaccharide, freeze dried in a stabilized matrix. RSE/CSE RATIO: The potency of this standard in Endotoxin units, (EU) has been determined to be \_EU/ng by the method formerly described in Appendix C (Gel-clot Technique) of the GUIDELINE ON VALIDATION OF THE LIMULUS AMEBOCYTE LYSATE TEST AS AN END PRODUCT ENDOTOXIN TEST FOR HUMAN AND ANIMAL PARENTERAL DRUGS, BIOLOGICAL PRODUCTS, AND MEDICAL DEVICES, published by the U.S. Food and Drug Administration. EX03492 CSE Exp. Date: \_ Dec 2023 LAL Reagent Lot: \_\_\_ M3021X LAL Exp. Date: Oct 2023 H0K354 \_ RSE/CSE Ratio: \_ 6 \_\_\_EU/ng Vial contents: \_\_\_\_60 \_\_ EU/vial RSE Lot: 0.25 \_\_EU/mL Geometric Mean Sensitivity with RSE: \_\_\_ IS/CSE RATIO: The Expert Committee on Biological Standardization of WHO has assigned a potency of the IS as 10,000 IU per vial of IS, so that 1 IU = 1 EU. The potency of this endotoxin standard in International (Endotoxin) Units, IU, has been designated as \_\_\_\_\_6\_\_\_IU/ng. **DIRECTIONS FOR USE:** Reconstitute the lyophilized material with \_\_\_\_\_\_ 3.0 \_\_\_mL of LAL reagent grade water to obtain 20 EU/mL or 20 IU/mL. Vortex mix vigorously for at least 5 minutes after rehydration, and for at least 1 minute immediately prior to each use. STORAGE: Store rehydrated material at 2-8°C for up to 4 weeks. Store lyophilized material at controlled room temperature or refrigerated as preferred. Diluted endotoxin should not be stored except under validated conditions. **CAUTION: DO NOT FREEZE ENDOTOXIN SOLUTIONS** Ouality Reviewed By: Oute: 15 Feb 2021

Date: 15 Feb 2021