

## **CERTIFICATE OF ANALYSIS**

**VIAL CONTENTS:** Endosafe® Control Standard Endotoxin is prepared from *E. coli* strain 055:B5. Each vial contains 10 ng of purified Lipopolysaccharide, freeze dried in a stabilized matrix.

RSE/CSE RATIO. The potency of this standard in Endotoxin units, (EO) has been determined to be
EU/ng by the method formerly described in Appendix C ( <i>Gel-clot Technique</i> ) of the GUIDELINE ON VALIDATION OF THE LIMULUS AMEBOCYTE LYSATE TEST AS AN END PRODUCT ENDOTOXIN TEST FOR HUMAN AND ANIMAL PARENTERAL DRUGS, BIOLOGICAL PRODUCTS, AND MEDICAL DEVICES, published by the U.S. Food and Drug Administration.
CSE Lot: <u>EX72123</u> CSE Exp. Date: <u>Aug 2020</u>
LAL Reagent Lot: J1601L LAL Exp. Date: Dec 2020
RSE Lot: H0K354 RSE/CSE Ratio: 10 EU/ng Vial contents: 100 EU/via
Geometric Mean Sensitivity with RSE: 0.125 EU/mL
IS/CSE RATIO: The Expert Committee on Biological Standardization of WHO has assigned a potency of the IS as 10,000 IU per vial of IS, so that 1 IU = 1 EU. The potency of this endotoxin standard in International (Endotoxin) Units, IU, has been designated asIU/ng.
<b>DIRECTIONS FOR USE:</b> Reconstitute the lyophilized material with5.0mL of LAL reagent grade water to obtain20EU/mL or20IU/mL. Vortex mix vigorously for at least 5 minutes after rehydration, and for at least 1 minute immediately prior to each use.
<b>STORAGE:</b> Store rehydrated material at 2-8°C for up to 4 weeks. Store lyophilized material at controlled room temperature or refrigerated as preferred. Diluted endotoxin should not be stored except under validated conditions.
CAUTION: DO NOT FREEZE ENDOTOXIN SOLUTIONS
Prepared By: Swallall Date: 07 Sep 2017
Prepared By: Date: 07 Sep 2017  Quality Reviewed By: Cynthu Rau Date: 07 Sep 2017

CA-GC10-05