

## CERTIFICATE OF ANALYSIS

VIAL CONTENTS: Endosafe® Control Standard Endotoxin is prepared from E. coli strain 055:B5. Each vial contains 500 ng of purified Lipopolysaccharide, freeze dried in a stabilized matrix. RSE/CSE RATIO: The potency of this standard in Endotoxin units, (EU) has been determined to be EU/ng by the method formerly described in Appendix C (Gel-clot Technique) of the GUIDELINE ON VALIDATION OF THE LIMULUS AMEBOCYTE LYSATE TEST AS AN END PRODUCT ENDOTOXIN TEST FOR HUMAN AND ANIMAL PARENTERAL DRUGS, BIOLOGICAL PRODUCTS, AND MEDICAL DEVICES, published by the U.S Food and Drug Administration. Jan 2022 CSE Lot: EM84083 CSE Exp. Date: \_ K4101X Jan 2021 LAL Exp. Date: LAL Reagent Lot: \_ EU/ng Vial contents: RSE Lot: RSE/CSE Ratio: \_ 0.125 EU/mL Geometric Mean Sensitivity with RSE: IS/CSE RATIO: The Expert Committee on Biological Standardization of WHO has assigned a potency of the IS as 10,000 IU per vial of IS, so that 1 IU = 1 EU. The potency of this endotoxin standard in International (Endotoxin) Units, IU, has been designated as \_\_\_\_\_10 IU/ng. DIRECTIONS FOR USE: Reconstitute the lyophilized material with \_\_\_\_\_\_ 5.0 \_\_\_\_ mL of LAL reagent grade water to obtain 1000 EU/mL or 1000 IU/mL. Vortex mix vigorously for at least 5 minutes after rehydration, and for at least 1 minute immediately prior to each use. STORAGE: Store rehydrated material at 2-8°C for up to 4 weeks. Store lyophilized material at controlled room temperature or refrigerated as preferred. Diluted endotoxin should not be stored except under validated conditions. **CAUTION: DO NOT FREEZE ENDOTOXIN SOLUTIONS** Date: 22 Mar 2018 Date: 23MAR2018 Quality Reviewed By:

CA-GC500-06